



primers. Based on their nucleotide sequences the 12 interacting clones were classified into 6 independent groups (see Table I).

TABLE I: Characterization of cDNA clones isolated by the yeast two hybrid screen.

Class	No. of clones	Identity	Mean RLU (Liquid assay)	Colour intensity (Filter assay)
1	6	Nedd4	2.86×10^6	++++
2	2	Htk	1.86×10^5	++
3	1	2.2412	5.18×10^6	++++
4	1	Proleosome	3.88×10^2	+/-
5	1	Solnatostatin	1.45×10^3	+/-
1	5	receptor		
6	1	L-arginine:glycine amidinotransferase	8.61×10^2	+/-

The 12 clones exhibiting activation of both the HIS 3 and lacZ reporter genes were divided into 6 groups by sequence analysis of their cDNA inserts. Results of β -galactosidase activity assays performed using two methodologies are shown. The liquid culture-derived method (Gallacto-Light TROPIX) is more quantitative; results are given in mean relative light units (RLU) and are normalized for the protein content of the samples. Blue/white screening of the cDNA clones was also performed using a colony lift filter assay (Clontech). The intensity of blue colour development over approximately 2h is scored from +/- (very weak) to ++++ (strong).

Six clones were partial cDNAs corresponding to Nedd4, a multidomain protein containing a calcium-dependent phospholipid binding (CaLB) domain, four WW domains and a C-terminal region homologous to the E6-AP carboxyl-terminus (Kumar *et al. Biochem. Biophys. Res. Commun.* 185, 1155-1161, 1992; Sudol *et al J. Biol. Chem.* 270, 14733-14741, 1995; Huijbregtse *et al Proc. Natl. Acad. Sci. USA* 92, 2563-2567, 1995). The latter is likely to confer E3 ubiquitin-protein ligase activity on Nedd4. The pACT2 clones isolated encoded the GalE domain together with the first 22 amino acids of the first WW domain.

RECEIVED
APR 08 2002
TECH CENTER 1600/2900